

Content available at: <https://www.ipinnovative.com/open-access-journals>

## International Journal of Pharmaceutical Chemistry and Analysis

Journal homepage: <https://www.ijpca.org/>

## Original Research Article

## Antibacterial efficacy of amla leaves

Vikas Sharma<sup>1,\*</sup>, Arti Heer<sup>1</sup>, Navneet Kour<sup>1</sup>, Shivangi Sharma<sup>2</sup><sup>1</sup>Division of Biochemistry, Faculty of Basic Sciences, Sher-e-Kashmir University of Agricultural Sciences and Technology of Jammu, Jammu and Kashmir, India<sup>2</sup>Dept. of Chemistry, University of Jammu, Jammu and Kashmir, India

## ARTICLE INFO

## Article history:

Received 10-08-2023

Accepted 08-09-2023

Available online 15-09-2023

## Keywords:

Amla leaves

Agar well diffusion method

Antibacterial

Methanolic extract

## ABSTRACT

The present study was carried out to evaluate *in vitro* antibacterial potential of methanolic extract from amla leaves via agar-well diffusion method against two gram positive (*Bacillus subtilis* MTCC2389, *Staphylococcus aureus* MTCC7443) and three gram negative (*Micrococcus luteus* MTCC4821, *Escherichia coli* MTCC2127, *Klebsiella pneumoniae* MTCC7162) strains. Susceptibility of plant extract was tested by serial microdilution method (MIC) and agar well diffusion method was determined. The extract showed maximum zone of inhibition against *Micrococcus luteus* (17.6 mm) followed by *Klebsiella pneumoniae* (16.5 mm) and *Bacillus subtilis* (16.2 mm). However, activity was also reported against *Staphylococcus aureus* (15.5 mm) and *Escherichia coli* (15 mm). The different concentrations of extract (10 $\mu$ l and 20 $\mu$ l) were used and the minimum inhibitory concentration (MIC) ranging from 62.5 to 2000  $\mu$ g/ml was compared with the standard drug Chloramphenicol (5 $\mu$ g). The MIC of methanolic extract was found 550  $\mu$ g/ml against *K. pneumoniae* and *B. subtilis*. The results of present study revealed that leaves of amla possesses antibacterial potential and source of new antibiotics. Therefore, medicinal plants are finding their way into pharmaceuticals, nutraceuticals and food Supplements that could be useful in chemotherapy to control infectious diseases.

This is an Open Access (OA) journal, and articles are distributed under the terms of the [Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License](https://creativecommons.org/licenses/by-nc-sa/4.0/), which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: [reprint@ipinnovative.com](mailto:reprint@ipinnovative.com)

## 1. Introduction

*Emblica officinalis* (amla/aonla) is a well known medicinally valued plant, the efficacy of amla fruit is widely proved, however, the use of its leaves is less investigated. Antimicrobial property of petroleum ether leaf extract of amla was tested by disc diffusion method and the MIC of antimicrobial activity, concentration ranging from 1000  $\mu$ g to 62.5  $\mu$ g was compared with the standard drug streptomycin (10  $\mu$ g) and amphotericin B (20  $\mu$ g). The study revealed that the extract has potential antibacterial activity, leaves have broad spectrum of antimicrobial activity and a potential source for new class of antibiotics

that could be useful in chemotherapy / control human infectious diseases (Elangovan et al., 2015). The aqueous crude extract of amla was screened by using agar well diffusion method against five human bacterial pathogens such as *Bacillus sp.*, *Lactobacillus sp.*, *Pseudomonas sp.*, *Proteus sp.* and *Streptococcus sp.* at the concentrations of 30, 60 and 90 $\mu$ l. The data clearly shows that the extract possesses strong inhibitory action against all the test bacterial pathogens (Kanthimathi and Soranam, 2013). *In vitro* antibacterial activity of aqueous, ethanolic and acetone extracts of fruit was evaluated against gram-positive vs. gram-negative bacteria employing *S. aureus* and *E. coli* respectively. All the extracts exhibited significant antibacterial activity, more against *S. aureus* than *E. coli*. Among the extracts, acetone inhibited the growth of *S.*

\* Corresponding author.

E-mail address: [vikas.skuast@gmail.com](mailto:vikas.skuast@gmail.com) (V. Sharma).

*aureus* and *E. coli* at minimum concentration of the extract (0.1  $\mu\text{g}$  and 1.0  $\mu\text{g}$  respectively). MIC for ethanol and aqueous extracts were 0.3 and 1.0  $\mu\text{g}$ , 1.5 and 3.75  $\mu\text{g}$ , for *S. aureus*, *E. coli* respectively.<sup>1–3</sup> It is concluded that *E. officinalis* is more inhibitory to gram-positive than the gram-negative bacteria (Varghese et al., 2013).

Extracts of different solvents of the leaf and bark of the plant were used to identify the bioactive compounds and antimicrobial activity was checked against different human pathogens (MTCC). Methanolic extract of amla showed highest zone of inhibition against *E. aerogens* and *E. faecalis* and acetate extract showed antifungal activity against *Rhizomucor* species (Sukanya et al., 2013). Antifungal property of *E. officinalis* was also reported against *Aspergillus* species (Satish et al., 2007). Ethanol and acetone extracts of fruit showed moderate activity against *F. equiseti* and *C. albicans* where griseofulvin was used as standard antibiotic (Hossain et al., 2012).<sup>4–8</sup>

Amla is also reported to possess potent free radical scavenging, antioxidant, anti-inflammatory, anti-mutagenic and immunomodulatory activities which are efficacious in the prevention and treatment of various diseases like cancer, atherosclerosis, diabetes, liver and heart diseases. Therefore, an attempt was made to assess the antibacterial property of *E. officinalis* using Agar-well diffusion method against some common bacterial pathogens.

In view of its medicinal and antimicrobial properties the present study aimed at assaying quantitatively the antibacterial activity of methanolic extracts of *E. officinalis* against gram positive and gram negative bacterial strains

## 2. Materials and Methods

### 2.1. Antimicrobial activity

#### 2.1.1. Determination of antibacterial activity

Qualitative analysis for evaluating antimicrobial activity of test material was carried out by agar-well diffusion method (Feyza et al., 2009) with modification. Two gram positive (*Bacillus subtilis*, MTCCC 2389 and *Staphylococcus aureus* MTCC7443) and three gram negative (*Micrococcus luteus*, MTCC4821., *Escherichia coli*, MTCC2127., *Klebsiella pneumoniae*, MTCC7162) bacterial strains were used in the present study. 20 mL of sterilized nutrient agar was inoculated with 100  $\mu\text{L}$  of bacterial suspension ( $10^8\text{CFU/mL}$ ) and then poured on to sterilized petri plates. The agar plates were left to solidify at room temperature.<sup>9–12</sup> A well of 6 mm diameter was aseptically bored into the agar plates and 20  $\mu\text{L}$  of the essential oil (diluted with DMSO, 1:1) was added in each well. Chloramphenicol (5 $\mu\text{g}$ ) was used as a positive reference to determine the sensitivity of bacteria and DMSO as negative reference. The plates were kept at 4 °C for 2 h to allow the dispersion and then incubated at 37 °C for 24 h.

### 2.2. Determination of MIC by broth dilution method

Broth dilution technique was used to determine the minimum inhibitory concentration of the test material against bacterial strains. One millilitre of nutrient broth was kept in each tube and autoclaved. The extract diluted with DMSO (1:1) was filtered with 0.22  $\mu\text{m}$  filter disk before use and then added to each tube to keep the final concentration ranging from 62.5  $\mu\text{g/mL}$ –2000  $\mu\text{g/mL}$ . The test bacterial suspension was added into each tube to yield bacterial density of  $10^6\text{CFU/mL}$  and the inoculated tubes were incubated at 37 °C for 24 h. Tubes containing nutrient broth without essential oil served as positive control, whereas those without bacteria as negative control. After incubation, 50 $\mu\text{L}$  of 0.2 mg/mL p-iodonitrotetrazolium violet (INT) was added in each tube to indicate the bacterial growth. The tubes were again incubated for 30 min at 37 °C. Development of pink colour in the tube (due to reduction of dye) indicated the bacterial growth whereas tubes without colour indicated no active bacterial growth. The lowest concentration at which no bacterial growth was observed (as indicated by colour) corresponded to the minimum inhibitory concentration (MIC). All the assays were performed in triplicate.

## 3. Results and Discussion


The explore for antimicrobials from natural resources has been received much attention to identify compounds that can act as suitable antimicrobials agent to replace synthetic products. Phytochemicals derived from plant products serve as a prototype to develop less toxic and more effective medicines in controlling the growth of microorganism (Kelmansone et al., 2000) and (Ahmad et al., 2001). These compounds have significant therapeutic application against human pathogens including bacteria, fungi or virus. Numerous studies have been conducted with the extracts of various plants, screening antimicrobial activity as well as for the discovery of new antimicrobial compounds (Guleria et al., 2006) and (Zakaria et al., 2007). The present study was carried out to evaluate in vitro antibacterial potential of methanolic extract of *Emblica officinalis* leaves. The antibacterial activity was screened by using Agar-well diffusion method against two Gram positive strains viz., *Bacillus subtilis* MTCC2389, *Staphylococcus aureus* MTCC7443 and Gram negative strains like *Escherischia coli* MTCC212 and *Klebsiella pneumoniae* MTCC7162 and *Micrococcus luteus* MTCC4821.

The extract showed maximum zone of inhibition against *Micrococcus luteus* (17.6mm) followed by *Klebsiella pneumoniae* (16.5mm) and *Bacillus subtilis* (16.2 mm) and lowest against *Staphylococcus aureus* (15.5mm) and *Escherischia coli* (15mm) (Table 1). The present study also revealed that the methanolic extracts of amla exhibited significant antibacterial activity more against

**Table 1:** Antibacterial analysis of *Emblica officinalis* leaves

Extract	Conc. (mg/mL)	Bacterial strains				
		B. subtilis MTCC 2389	M. luteus MTCC 4821	S. aureus MTCC 7443	E. coli MTCC 2127	K. pneumoniae MTCC 7162
Methanolic	100 mg/mL (10 $\mu$ l)					
		Zone of inhibition (mm)				
MIC	62.5 $\mu$ g/ml-2000 $\mu$ g/ml	16.3 $\pm$ 0.15 500	17.6 $\pm$ 0.20 500	15 $\pm$ 0.1 500	15.2 $\pm$ 0.15 500	16.3 $\pm$ 0.16 250
Positive control CMP <sup>+</sup>	1 mg/mL (10 $\mu$ l)	36 $\pm$ 1.3	35.6 $\pm$ 1.2	26.2 $\pm$ 0.9	35 $\pm$ 1.1	35 $\pm$ 1.3
Negative control DMSO	10 $\mu$ l	-	-	-	-	-

Values indicating maximum zone of inhibition of test component are given in bold numbers CMP<sup>+</sup>: Chloramphenicol; DMSO: Dimethylsulfoxide Mark (-) indicates no activity



**Fig. 1:** Antibacterial activity of *Emblica officinalis* extract against **A:** *Klebsiella pneumoniae* and **B:** *Micrococcus luteus* by agar well diffusion assay **C:** *Staphylococcus aureus* and **D:** *Bacillus subtilis* by agar well diffusion assay

*Micrococcus luteus* than others bacterial strains. The different concentrations of extract were used for these tests were 10 $\mu$ l and 20 $\mu$ l. The minimum inhibitory concentration (MIC) of antimicrobial activity of the amla extract at a concentration ranging from 2000  $\mu$ g to 62.5  $\mu$ g was compared with the standard drug Chloramphenicol (disc 5 $\mu$ g). The antibacterial activity exhibited by the *E. officinalis* could be attributed to the presence of bioactive components namely flavonoids, phenols, saponins, tannins in the plant Javale and Sabnis (2010). The results of present study revealed that leaves of amla contain high antibacterial potential and source of new classes of antibiotics that could be useful in chemotherapy and control on human infectious diseases.

#### 4. Conclusion

The present study disclosed the importance of natural medicinal plant extract to control pathogenic bacteria which pose threat to human health and it also concludes that leaves of amla contain high antimicrobial properties. The presence of phytochemicals like flavonoids, tannins, saponins glycosides and phenolics compounds are major constituents in *Emblica officinalis* which may acknowledge the medicinal property of this plant. This implies that the methanolic extract may indeed be effective in the management of diseases caused by these encountered pathogens and supporting its ethno medicinal uses and this plant also safe, potent and cost effective to treat many infectious diseases of livestock, poultry and human. There is need for further investigation of this plant in order to identify and isolate its antimicrobial agent.

#### 5. Source of Funding

None.

#### 6. Conflict of Interest

None.

#### References

1. Sukanya MK, Aruna SS. Phytochemical analysis, antimicrobial screening and antihelminthic properties of *Phyllanthus emblica*. *Int J Pharma Bio Sci.* 2013;4(4):55–64.
2. Varghese LS, Ninan MA, N A, S S, S J. Comparative antibacterial activity of fruit extracts of *Emblica officinalis* Gaertn against gram-positive versus gram-negative bacteria. *Biomedical and Pharmacology Journal.* 2013;6:447–452.
3. Zakaria Z, Sreenivasan S, Mohamad M. Antimicrobial Activity of *Piper ribesoides* Root Extract against *Staphylococcus aureus*. *J App Biol Sci.* 2007;1(3):87–90.
4. Beg AI. Antimicrobial and phytochemical studies on 45 Indian medicinal plants against multiple drug resistant human pathogens. *J Ethnopharma.* 2001;74(2):113–23.

5. Elangovan NM, Dhanarajan MS, Elangovan I. Evaluation of antibacterial and antifungal activity of *Phyllanthus emblica* leaf extract. *Int Res J Pharma Biosci.* 2015;2(2):59–66.
6. Feyza O, Belma A, Sahlan O, Senol A. Essential oil composition of antimicrobial and antioxidant activity of *Satureja cuneifolia* Ten. *Food Chem.* 2009;112:874–9.
7. Guleria S, Kumar A. Antifungal activity of some Himalayan medicinal plants using direct bioautography. *J Cell Mol Bio.* 2006;5:95–8.
8. Hossain MM, Mazumder K, Hossen SMM, Tanmy TT, Rashid MDJ. In vitro studies on antibacterial and antifungal activities of *Emblica officinalis*. *Int J Pharm Sci Res.* 2012;3:1124–7.
9. Sabnis J. Antimicrobial properties and phytochemical analysis of *Emblica officinalis*. *Asian J Exp Biol Sci.* 2010;31(3):91–5.
10. Soranam KM. Antibacterial effects of *Emblica officinalis* and *Phyllanthus niruri* crude extracts against bacterial pathogen. *Int J Pharm Clin Sci.* 2013;3(3):20–3.
11. Kelmanson JE, Jager AK, Staden V. Zulu medicinal plants with antibacterial activity. *J Ethnopharmacol.* 2000;69(3):241–6.
12. Satish S, Mohana DC, Raghavendra MP, Raveesha KA. Antifungal activity of some plant extracts against important seed borne pathogens

of *Aspergillus* sp. *J Agricul Technol.* 2007;3:109–19.

### Author biography

**Vikas Sharma**, Associate Professor

**Arti Heer**, Assistant Professor

**Navneet Kour**, –

**Shivangi Sharma**, –

**Cite this article:** Sharma V, Heer A, Kour N, Sharma S. Antibacterial efficacy of amla leaves. *Int J Pharm Chem Anal* 2023;10(3):205-208.