

Content available at: <https://www.ipinnovative.com/open-access-journals>

International Journal of Pharmaceutical Chemistry and Analysis

Journal homepage: <https://www.ijpca.org/>

Original Research Article

Study the effect of Vitamin D3 in newly diagnosed acute myeloid leukemia

Shatha M. J. Al-Khateeb¹, Raghada Shams Akram¹, Mohammad Ali Kareem², Fatma Abdalhamza Obed^{3*}¹Dept. of Pharmaceutical Chemistry, College of Medicine, Mustansiriyah University, Baghdad, Iraq²Ministry of Health Baghdad Health, Baghdad, Iraq³National Center of Hematology, Mustansiriyah University, Baghdad, Iraq

ARTICLE INFO

Article history:

Received 12-12-2023

Accepted 26-12-2023

Available online 26-03-2024

Keywords:

Acute myeloid leukemia

Vitamin D3

Ferritin

Male

Female

ABSTRACT

Objective: This study is aimed to put light on the effect of vitamin D3 (VitD3) in newly diagnosed acute myeloid leukemia (AML) patients.**Materials and Methods:** The AML patients consisted of sixty patients (30 male, 30 female). The average age of patients was (36.44±8.44) years (range 25–45). Thirty healthy subjects were considered as controls (15 male and 15 female), and the average age of control group was (32.22±5.13) years. All subjects had to go through clinical examination to determine existence of other diseases in addition to evaluate some tests as complete blood count [hemoglobin (Hb), platelet (PLT), and white blood cells (WBCs)], ferritin, FBS (fasting blood sugar), TG (triglycerides), TC (total cholesterol), and HDL (high density lipoprotein cholesterol) (HDL-Ch), urea, and creatinine. Vitamin D3 was measured by minividase Biomerux/French.**Results:** The VitD3 was highly significantly lower in AML patients than in controls (p=0.001). While, VitD3 was lower in male than in female AML patients but the difference was not statistically significant. There were statistically significant positive correlations between VitD3 and Hb, PLT, Wbc, ferritin, FBS, and TG in both male and female AML patients.**Conclusion:** These findings suggest that lower serum concentrations of vitamin D3 may possibly be considered as a potential factor for early assessment of acute myeloid leukemia.This is an Open Access (OA) journal, and articles are distributed under the terms of the [Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License](https://creativecommons.org/licenses/by-nc-sa/4.0/), which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.For reprints contact: reprint@ipinnovative.com

1. Introduction

Acute myeloid leukemia (AML) which is defined as a hematological malignancy characterized by the proliferation of progranulocytes or myeloblasts that failed to developed normal myeloid differentiation.¹ It comprises a heterogeneous group of clinically and biologically aggressive disorders which driven by any combination or number of chromosomal abnormalities, recurrent gene mutations, and/or altered signaling pathways.² Based on the French-American-British leukemia experts, Acute myeloid

leukemia are morphologically classified into 8 subtypes(M0 to M7). Markedly, over 700 chromosomal abnormalities was detected in leukemic cells of the AML patients.³ In 2018, the global incidence of AML has been estimated to be nearly 130,000.⁴ AML tends to mainly affect older adults, with a median age of onset of sixty-eight years.⁵

Vitamin D (VitD), is a fat-soluble vitamin that is obtained from sun exposure, food, and supplements.⁶ VitD is a steroid hormone which produced by human skin as a result of ultraviolet (UV) radiation stimulus.⁷ VitD is important for calcium as well as bone metabolism. In addition, it has other roles that includes: reducing insulin resistance , ameliorating oxidative stress, and reducing the possibility

* Corresponding author.

E-mail address: fatma.hamza@gmail.com (F. A. Obed).

of prevalent metabolic syndrome.⁸

1,25-Dihydroxy-vitamin D₃, a vitamin D (cholecalciferol) active metabolite (1,25D₃, calcitriol), is a secosteroid hormone with a variety of biological functions.⁹ regulating the uptake and distribution of necessary minerals, such as calcium, magnesium, and phosphorus, which are essential for maintaining bone health, is one of 1,25D₃'s key physiological roles.¹⁰ VitD₃ regulates gene transcription through binding to vitamin D receptor (VDR). VitD₃, which is not dependent on transcription, can also elicit rapid intracellular responses in vitro.¹¹

VitD was implicated in the pathogenesis of hematologic malignancies and showed promise as an anticancer drug. Serum 25(OH) D₃ concentrations, the precursor of calcitriol, are usually lower in subjects with hematological disorder in comparison with healthy subjects. A worse illness outcome is commonly linked to this. Moreover, cancer models both in vitro and in vivo have shown that sick cells frequently produce high levels of the VDR, which is required for many of the anticancer effects. Importantly, "in abnormal hematological cells, vitamin D supplementation promotes apoptosis, induces differentiation, inhibits proliferation, sensitizes tumor cells to other anti-cancer therapies, and reduces the production of pro-inflammatory cytokines".¹²

This study is aimed to put light on the effect of VitD₃ in patients with newly diagnosed AML.

2. Subjects and Methods

Acute myeloid leukemia patients consisted of sixty patients (30 male, 30 female). The patients' average age was (36.44±8.44) years (range 25-45). Thirty healthy subjects were considered as controls (15 male and 15 female), while the control group's average age was (32.22±5.13) years. This investigation was done at the National Center of Hematology during the period from 1st December 2020 till 30th June 2021. All subjects had to go through clinical examination to determine existence of other diseases in addition to evaluate some tests as complete blood count [hemoglobin (Hb), platelet (PLT), and white blood cells (WBCs)], ferritin, urea, creatinine, Fasting blood sugar, total cholesterol, triglycerides, and high density lipoprotein (HDL) (FBS). Minividase Biomerux/French was used to calculate vitamin D₃ levels.

In accordance with the Helsinki Ethical Guidelines, this investigation was completed. Using SPSS, a statistical analysis was performed. The investigation's findings are expressed as mean ± standard deviation. Using the student's t-test, the mean values of the two groups were compared. The P-value 0.05 or lower was regarded as significant.

3. Results

The hematological and biochemical parameters of the AML patients and controls are summarized in Table 1. FBS, TC, TG, and urea were all statistically significantly higher in newly diagnosed AML patients compared to controls (all $p=0.05$), and WBC ($p=0.01$). In comparison to controls, PLT and ferritin levels in AML patients were highly significantly lower. Compared to the controls, VitD₃ levels in AML patients were significantly lower ($p=0.001$). Hb and HDL-Ch were statistically significantly decrease ($p=0.05$ and $p=0.016$, respectively) in AML patients compared to controls. However, regarding creatinine levels, there were no statistically significant changes between AML patients and controls.

The AML patients are broken down by sex in Table 2. Male versus female AML patients had statistically significantly higher levels of Hb, PLT, TC, TG, and ferritin. Although the difference was not statistically significant, male AML patients had lower VitD₃ levels than female AML patients. The WBC was statistically significantly lower in male compared with female AML patients ($P=0.01$). Nevertheless, There were no statistically significant differences found in FBS, HDL-Ch, urea, and creatinine between male and female AML patients.

Table 3 showed that there were statistically significant positive correlations between VitD₃ and Hb ($r=0.333$, $r=0.244$), PLT ($r=0.330$, $r=0.296$), Wbc ($r=0.305$, $r=0.284$), ferritin ($r=0.338$, $r=0.374$), FBS ($r=0.325$, $r=0.203$), and TG ($r=0.662$, $r=0.665$) in male and female AML patients respectively.

4. Discussion

In this investigation, highly significantly lower VitD₃ level was found in AML subjects compared to the control group. Similar to earlier investigations.¹³⁻¹⁶ One of these investigations was performed by Bobilev et al.¹⁴ who detected low levels of 1.25 (OH)D in human acute myeloid leukemia cell lines.

In addition, Elkerdany et al.¹³ and Zidan et al.¹⁵ established that VitD contribute to the regulation of cellular proliferation, differentiation accompanied by angiogenesis and apoptosis. Furthermore, Munker and his team¹⁷ revealed that VitD is a highly potent inhibitor of the CD34 leukemic cells, whereas maintaining activity of the normal CD34 hematopoietic stem-progenitor cells. Lappe et al. and these findings are usually in agreement.¹⁸ They demonstrated that a major independent predictor of the risk of cancer was the serum 25(OH)D level., in addition to vitamin D and supplemented calcium being associated with a lower risk of cancer.. Additionally, Thomas and his coworkers¹⁹ revealed a significant relationship between malignant cell burden and circulatory 25(OH)D. They reported that the lower concentrations of

Table 1: The hematological and biochemical parameters of the aml patients and controls

Parameters	AML N (60)	Control N (30)	P-value
	Mean±SD	Mean±SD	
Hb (g/dl)	10.6±1.66	13.7±1.12	0.05*
PLT(103/ μ l)	98.5±7.7	206.2±18.1	0.001**
WBCs (103/ μ l)	25.5±1.55	8.44±1.23	0.01**
Ferritin (ng/ml)	35.4±5.56	85.1±15.5	0.01**
FBS (mg/dl)	97.4±12.5	79.9±13.32	0.05*
TC (mg/dl)	220.11±22.41	168.1±18.02	0.05*
TG (mg/dl)	168.33±13.4	120.5±11.4	0.05*
HDL-Ch (mg/dl)	44.5±4.01	46.1±5.8	0.016*
Urea (mg/dl)	45.4±6.06	25.3±5.2	0.05*
Creatinine (mg/dl)	0.83±0.31	0.65±0.11	0.220
Vitamin D3 (ng/ml)	12.4±6.81	38.6±4.2	0.001**

Levels of significance= *P<0.05, and **P<0.01.
Hb: hemoglobin, PLT: platelet, WBCs: white blood cells, FBS: fasting blood sugar, TC: total cholesterol, TG: triglyceride, HDL-Ch: high density lipoprotein cholesterol.

Table 2: The hematological and biochemical parameters between male and female aml patients

Parameters	Male N (30)	Female N (30)	P-value
	Mean±SD	Mean±SD	
Hb (g/dl)	12.0±1.50	9.32±0.88	0.05*
PLT (103/ μ l)	69.5±5.71	35.10±5.11	0.05*
WBCs (103/ μ l)	10.8±1.66	16.6±0.74	0.01**
Ferritin (ng/ml)	30.7±10.0	15.2±8.12	0.01**
FBS (mg/dl)	87.3±8.77	94.2±8.4	0.130
TC (mg/dl)	224±15.41	198.4±31.22	0.05*
TG (mg/dl)	187.5±11.0	170.44±10.14	0.05*
HDL-Ch (mg/dl)	44.9±10.4	46.2±5.21	0.132
Urea (mg/dl)	25.4±1.7	26.3±1.8	0.142
Creatinine (mg/dl)	0.73 ±0.32	0.72±0.43	0.175
Vitamin D3 (ng/ml)	16.86±3.40	28.9±3.01	0.51

Levels of significance= *P<0.05, and **P<0.01.
Hb: hemoglobin, PLT: platelet, WBCs: white blood cells, FBS: fasting blood sugar, TC: total cholesterol, TG: triglyceride, HDL-Ch: high density lipoprotein cholesterol.

Table 3: Correlation coefficient between VitD3 levels and all parameters in both genders (male and female) AML patients.

Parameters	Vitamin D3	
	Male	Female
	r	r
Hb (g/dl)	0.333*	0.244*
PLT (103/ μ l)	0.330*	0.296*
WBCs (103/ μ l)	0.305*	0.284*
Ferritin (ng/ml)	0.338*	0.374*
FBS (mg/dl)	0.325*	0.203*
TC (mg/dl)	0.139	0.026
TG (mg/dl)	0.662**	0.665**
HDL-Ch (mg/dl)	0.149	0.113
Urea (mg/dl)	0.131	0.102
Creatinine (mg/dl)	0.102	0.032

Levels of significance= *P<0.05, and **P<0.01.

circulatory 25(OH)D seemed to be associated with disease aggressiveness, the progressive phase of the disease along with poor response to treatment. Hence, it is a potential biomarker of prognosis in subjects with acute myeloid leukemia.

In this investigation, no statistically differences in VitD3 levels were found between male and female AML patients. Epidemiologic studies implied a relationship between AML and low 25(OH) vitamin D3 concentrations. For instance, a research in United Arab Emirates²⁰ reported female predominance in AML, even though the population of United Arab Emirates consists of more males than females, and even though it is very widely known that acute myeloid leukemia is more common in males. These results indicate that the low VitD3 concentrations secondary to the practice of females wearing conservative clothing can possibly contribute to higher incidence of acute myeloid leukemia.²⁰

In this study, VitD3 levels showed significant positive correlations with Hb, PLT and WBC in both male and female AML groups. Elkerdany et al.¹³ reported a positive correlation between VitD and PLT, while no correlations between VitD with Hb and WBC were found in AML subjects.

Globally, numerous investigations were performed to examine the relationship between ferritin and VitD. Serum ferritin concentrations are regulated by hepcidin, that plays a central role in decreasing human intestinal iron absorption. VitD “regulates the hepcidin-ferroportin axis in macrophages”, in addition, the increase of VitD is known to lower systemic concentrations of hepcidin which ameliorate anemia.²¹ In this investigation, VitD3 and serum ferritin levels showed a positive correlation. This result agreed with other investigations, where a positive association was seen in adults,²² children and adolescents,²³ and fibroid uterus.²⁴

It has been reported in different investigations that VitD has an important role in control of circulating glucose concentrations and its utilization in target tissues. Numerous human and animal investigations have demonstrated the role of VitD in insulin synthesis in addition to its secretion from the beta cells.²⁵ It has shown that there was a significant positive correlation between VitD3 and FBS in male and female AML patients in the current study. Nonetheless, this result contradicts those of other previous studies in different populations.^{26,27} Many lines of strong evidence suggest that vitamin D3 affects insulin resistance, systemic inflammation, and impaired pancreatic beta cell activity.²⁸ VitD3 can alter insulin sensitivity to raise blood glucose levels because it has been found that numerous tissues, including pancreatic islet beta cells, contain VitD3 receptors.²⁶

Due to the lack of sufficient sun exposure, a main cause of VitD deficiency, is considered to be linked to elevated serum cholesterol concentrations,²⁹ it seems plausible that VitD deficiency plays an essential role in

hypercholesterolemia. Moreover, the association between cholesterol and VitD status might be evaluated since both serum cholesterol (involving TC, HDL-Ch, and LDL-Ch) and 25(OH)D3 concentrations are routinely assessed from the blood samples that taken in research investigations. A literature search showed a common trend in which low 25(OH)D3 concentrations were linked to increased LDL-Ch and TC concentrations, increased LDL-Ch/HDL-Ch ratio and TC/HDL-Ch ratio, or reduced HDL-Ch concentrations. These trends have held true in diverse patient populations which included individuals of different ages and ethnicities. Consequently, there is significant evidence to support the idea that VitD deficiency plays an essential role in hypercholesterolemia. Nevertheless, these “cross-sectional” studies might only offer relationships and not causality.³⁰ The presence of cholesterol in the blood and tissues has been consistently demonstrated to have a major role in the pathogenesis of ischemic heart disease, but the relationship of cholesterol to cancer, such as breast, leukemia, and colorectal cancer, has also been revealed. In cell membranes, the cholesterol content is tightly regulated, furthermore, this process of regulation includes the uptake of LDL-Ch. Nonetheless, intriguingly, cholesterol accumulation was detected in various types of solid tumours, particularly prostate and oral cancers. Moreover, dysregulated cholesterol metabolism represents a significant metabolic alteration in several malignancies, involving breast and lung cancers as well as myeloid leukemia.³¹ Males with hypertriglyceridemia may benefit from lowered serum TG concentration as well as extensive colonoscopic surveillance in colorectal cancer, even if a high serum TG concentration does not appear to be mechanically implicated in the formation of most malignancies.³² A relationship between colorectal cancer and high serum TG was as well detected.³³ In this investigation, VitD3 and TG showed a statistically significant positive correlation in both male and female AML patients.

5. Conclusion

the result of this investigation revealed that lower serum concentrations of vitamin D3 may possibly be considered as a potential factor for assessment of acute myeloid leukemia.

6. Conflict of Interest

None.

7. Source of Funding

None.


8. Acknowledgment


The authors wish to acknowledge support from Mustansiriyah University (www.uomustansiriyah.edu.iq), Baghdad, Iraq.

References


- Obed FA, Shafeeq NK, Akram RS, Hmeed EZ. Study the Effect of High Sensitive-C reactive protein in newly Diagnosed Acute Myeloid Leukemia with other Hematological markers. *Res J Pharm Technol*. 2023;16(1):119–41.
- Aitken M, Ravandi F, Patel KP, Short NJ. Prognostic and therapeutic implications of measurable residual disease in acute myeloid leukemia. *J Hematol Oncol*. 2021;14(1):1–15.
- Pan P, Chen X. Nuclear receptors as potential therapeutic targets for myeloid leukemia. *Cells*. 2020;9(9):1–16.
- Bray F, Ferlay J, Soerjomataram I, Siegel RL, Torre LA, Jemal A. Global cancer statistics 2018: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA Cancer J Clin*. 2018;68(6):394–424.
- Forsythe A, Sandman K. What does the economic burden of acute myeloid leukemia treatment look like for the next decade? An analysis of key findings, challenges and recommendations. *J Blood Med*. 2021;12:245–55.
- Tahir N, Akram RS, Al-Rubae'i SHN, Nsaif AS. Vitamin D3 status and insulin resistance in Iraqi patients with osteoarthritis disease in Baghdad governorate-Iraq. *AIP Conference Proceedings*.
- Ishikawa L, Colavite PM, Fraga-Silva T, Mimura L, França T, Zorzella-Pezavento S. Vitamin D Deficiency and Rheumatoid Arthritis. *Clin Rev Allergy Immunol*. 2017;52(3):373–88.
- Chen H, Zhang H, Xie H, Zheng J, Lin M, Chen J. Maternal, umbilical arterial metabolic levels and placental Nrf2/CBR1 expression in pregnancies with and without 25-hydroxyvitamin D deficiency. *Gynecol Endocrinol*. 2021;37(9):807–20.
- Norman AW. The history of the discovery of vitamin d and its daughter steroid hormone. *Ann Nutr Metab*. 2012;61(3):199–206.
- Lieben L, Carmeliet G. Vitamin D signaling in osteocytes: Effects on bone and mineral homeostasis. *Bone [Internet]*. 2013;54(2):237–80.
- Kutner A, Brown G. Vitamins D: Relationship between structure and biological activity. *Int J Mol Sci*. 2018;19(7):1–11.
- Kulling PM, Olson KC, Olson TL, Feith DJ, Loughran TP. Vitamin D in hematological disorders and malignancies. *Eur J Haematol*. 2017;98(3):187–97.
- Eissa D, Moussa M, Elkerdany T. Serum 25-hydroxyvitamin D levels in relation to disease status and prognosis in acute myeloid leukemia. *Egypt J Haematol*. 2014;39(2):47.
- Bobilev I, Novik V, Levi I, Shpilberg O, Levy J, Sharon Y. The Nrf2 transcription factor is a positive regulator of myeloid differentiation of acute myeloid leukemia cells. *Cancer Biol Ther*. 2011;11(3):317–29.
- Zidan A, Omran A, Ghonaim R, Abu-Taleb F, Elgohary T. The Prognostic Value of Vitamin D Insufficiency & Vitamin D Receptor Gene Polymorphism in Adult Acute Myeloid Leukemia Patients. *J Cancer Ther*. 2019;10(5):361–70.
- Ismael AA. Biological markers study of acute and chronic myeloid leukemia. *Middle East J Intern Med*. 2014;7(2):3–8.
- Munker R, Norman A, Koefler HP. Effect on clonal proliferation and differentiation of human myeloid cells. *J Clin Invest*. 1986;78(2):424–54.
- Lappe JM, Travers-Gustafson D, Davies KM, Recker RR, Heaney RP. Vitamin D and calcium supplementation reduces cancer risk: Results of a randomized trial. *Am J Clin Nutr*. 2007;85(3):794.
- Thomas X, Chelghoum Y, Fanari N, Cannas G. Serum 25- hydroxyvitamin D levels are associated with prognosis in hematological malignancies. *Hematology*. 2011;16(5):278–83.
- Hassan IB, Islam SI, Alizadeh H, Kristensen J, Kambal A, Sondag S, et al. Acute leukemia among the adult population of United Arab Emirates: An epidemiological study. *Leuk Lymphoma*. 2009;50(7):1138–85.
- Kumari S, Singh S. Association of vitamin D, calcium and phosphate with uterine fibroid in premenopausal women of coastal Odisha. *Int J Sci Res*. 2019;8(2):27–36.
- Seong JM, Yoon YG, Lee KS, Bae NY, Gi MY, Yoon H. Gender difference in relationship between serum ferritin and 25-hydroxyvitamin D in Korean adults. *PLoS One*. 2017;12(5):1–13.
- Andiran N, Çelik N, Akça H, Doğan G. Vitamin D deficiency in children and adolescents. *JCRPE J Clin Res Pediatr Endocrinol*. 2012;4(1):25–34.
- Kumari S, Swetha P, Krishnan RS, Nayak S, Singh S. The Association Between Ferritin and Vitamin D Levels in Premenopausal Fibroid Uterus Cases With Anemia. *Cureus*. 2021;13(2):e13392. doi:10.7759/cureus.13392.
- Alvarez JA, Ashraf A. Role of vitamin D in insulin secretion and insulin sensitivity for glucose homeostasis. *Int J Endocrinol*. 2009;p. 351385. doi:10.1155/2010/351385.
- Kanakaraju K, Ranganathan RS, Shankar R. Correlation of Vitamin D3 Levels and the Blood Sugar Parameters among the Patients with type 2 Diabetes Mellitus. *Int J Contemp Med Res*. 2017;4(4):844–51.
- O'hartaigh B, Thomas N, Silbernagel G, Bosch G, Pilz JA, Loerbroks S, et al. Association of 25-hydroxyvitamin D with type 2 diabetes among patients undergoing coronary angiography: Cross-sectional findings from the Ludwigshafen Risk and Cardiovascular Health (LURIC) Study. *Clin Endocrinol (Oxf)*. 2013;79(2):192–200.
- Husemoen L, Thuesen BH, Fenger M, Jorgensen T, Glümer C, Svensson J. Serum 25(OH)D and type 2 diabetes association in a general population: A prospective study. *Diab Care*. 2012;35(8):1695–700.
- Walker AR, Shor A. Sunlight, cholesterol and coronary heart disease. *QJM*. 1997;90(2):153–7.
- Quach HP. Vitamin D Receptor (VDR) and Cholesterol Homeostasis : Interplay of VDR Enzyme Targets and Vitamin D Deficiency by Vitamin D Receptor (VDR) and Cholesterol Homeostasis : Interplay of VDR Enzyme Targets and Vitamin D Deficiency. *Best Pract Res Clin Endocrinol Metab*. 2016;25(4):543–59.
- Akram RS, Isaa MQ, Mohammed AA. Fetuin-A levels and Insulin Resistance in Obese and Non-Obese Iraqi Children. *J Popul Ther Clin Pharmacol*. 2023;30(6):414–34.
- Tabuchi M, Kitayama J, Nagawa H. Hypertriglyceridemia is positively correlated with the development of colorectal tubular adenoma in Japanese men. *World J Gastroenterol*. 2006;12(8):1261–5.
- Eyssen GM. Epidemiology of Colorectal Cancer Revisited: Are Serum Triglycerides and/or Plasma Glucose Associated with Risk? *Cancer Epidemiol Biomarkers Prev*. 1994;3(8):687–95.

Author biography

Shatha M. J. Al-Khateeb, Assistant Professor  <https://orcid.org/0000-0003-1872-515X>

Raghada Shams Akram, Assistant Lecturer  <https://orcid.org/0000-0002-9500-6485>

Mohammad Ali Kareem, Lecturer  <https://orcid.org/0009-0005-9423-1209>

Fatma Abdalhamza Obed, Epidemiology Specialist  <https://orcid.org/0000-0003-4864-0763>

Cite this article: Shatha M. J. Al-Khateeb, Akram RS, Ali Kareem M, Obed FA. Study the effect of Vitamin D3 in newly diagnosed acute myeloid leukemia. *Int J Pharm Chem Anal* 2024;11(1):72-76.