

Content available at: <https://www.ipinnovative.com/open-access-journals>

International Journal of Pharmaceutical Chemistry and Analysis

Journal homepage: <https://www.ijpca.org/>

Review Article

The intersection of pharmacology and dentistry: A comprehensive overview of pharmacotherapeutics

Shailesh Shenoy¹, Zameer Pasha², Suresh Rajendiran³, Syed Owais Ahmed⁴,
Krishan Kant⁵, Akshaya Narayan Shetti^{6,*}

¹Dept. of Pediatric and Preventive Dentistry, Yenepoya Dental College, Yenepoya University, Mangaluru, Karnataka, India

²Durrat Al-Alammi Dental Clinic, Al-Majmaah, Riyadh, Saudi Arabia

³Meenakshi Medical College Hospital and Research Institute, Kanchipuram. Faculty of Medicine, Meenakshi Academy of Higher Education & Research (MAHER) Deemed to be University, Chennai, Tamil Nadu, India

⁴Dept. of Periodontology and Implantology, Sri Siddhartha Dental College and Hospital, Tumakuru, Karnataka, India

⁵Dept. of Periodontology, Post Graduate Institute of Dental Science, Rohtak, Haryana, India

⁶Dept. of Anaesthesiology and Critical Care, Dr Balasaheb Vikhe Patil Rural Medical College, Loni, Maharashtra, India



ARTICLE INFO

Article history:

Received 12-04-2023

Accepted 10-05-2023

Available online 01-07-2023

Keywords:

Pharmacotherapy

Pharmacotherapeutics

Dentistry

Antibiotics

ABSTRACT

Pharmacotherapy is an essential component of dental practice to manage pain, prevent infections, and maintain oral health. Analgesics such as non steroidal anti-inflammatory drugs (NSAIDs) and opioids are commonly used to manage pain during and after dental procedures. Antibiotics are prescribed to treat and prevent infections, but their use should be limited to reduce the risk of antibiotic resistance. Local anesthetics provide safe and effective pain relief during dental procedures. Fluorides are used to prevent tooth decay and strengthen tooth enamel. Dental healthcare providers should evaluate each patient's individual needs and medical history to determine the most appropriate pharmacotherapy. Patient education on the proper use of medications and potential side effects is also important. By incorporating pharmacotherapy as part of a comprehensive dental care plan, patients can experience a more comfortable and successful dental experience.

This is an Open Access (OA) journal, and articles are distributed under the terms of the [Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License](https://creativecommons.org/licenses/by-nc-sa/4.0/), which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: reprint@ipinnovative.com

1. Introduction

Pharmacotherapeutics in dentistry refers to the use of medications to prevent, treat, or manage dental conditions and oral diseases. Dentists and dental healthcare providers commonly use medications to control pain, infection, inflammation, anxiety, and other conditions related to the mouth and teeth.¹ Some common medications used in dentistry include: Analgesics: These are pain-relieving medications that can be prescribed to manage dental pain. Examples include acetaminophen, non steroidal anti-inflammatory drugs (NSAIDs), and opioids.² Antibiotics:

These medications are used to treat bacterial infections such as periodontitis, dental abscesses, and necrotizing ulcerative gingivitis.³ Antifungal medications: These are used to treat oral thrush, a fungal infection that can occur in the mouth.⁴ Local anesthetics: These medications are used to numb a specific area of the mouth to make dental procedures more comfortable.⁵ Sedatives and anxiolytics: These medications are used to help patients relax during dental procedures, especially for those with dental anxiety.⁶ Fluoride: This mineral is used to strengthen teeth and prevent tooth decay. It can be applied topically or ingested through supplements. Chlorhexidine: This medication is used as a mouthwash to prevent and treat gum disease.

* Corresponding author.

E-mail address: aksnsdr@gmail.com (A. N. Shetti).

It's important to note that not all dental conditions require medication, and some may require referral to a specialist. Additionally, proper medication use and dosages should always be discussed with a dental healthcare provider or pharmacist.⁷

2. Role of Analgesics in Dentistry

The role of analgesics in dentistry is primarily to provide pain relief to patients experiencing dental pain. Pain is a common symptom associated with various dental conditions, such as tooth decay, gum disease, and dental trauma, among others. Pain can range from mild discomfort to severe, debilitating pain that can affect a person's daily activities and quality of life.⁸

Analgesics are medications that alleviate pain by blocking the transmission of pain signals to the brain or by reducing inflammation in the affected area. In dentistry, analgesics are commonly used to manage dental pain and discomfort associated with various procedures, such as tooth extraction, root canal therapy, and periodontal surgery, among others. There are different types of analgesics used in dentistry, including non-opioid analgesics, opioids, and local anesthetics. Non-opioid analgesics, such as acetaminophen and nonsteroidal anti-inflammatory drugs (NSAIDs), are commonly used for mild to moderate dental pain. They work by reducing inflammation and blocking the production of certain chemicals in the body that cause pain and fever.⁹ Opioids, such as codeine and hydrocodone, are stronger pain relievers that are prescribed for moderate to severe dental pain. They work by binding to specific receptors in the brain and spinal cord to reduce the perception of pain. However, opioids carry the risk of addiction, tolerance, and other adverse effects, and should only be used under close supervision by a healthcare provider. Local anesthetics, such as lidocaine and articaine, are used to numb the affected area during dental procedures to provide immediate pain relief. They work by blocking the transmission of pain signals from the nerves in the affected area to the brain. Proper use and dosage of analgesics in dentistry is essential to ensure that patients receive effective pain relief while minimizing the risk of adverse effects. Dental healthcare providers should assess the patient's pain level and medical history before prescribing analgesics and provide clear instructions on how to take them. Additionally, patients should be advised on the potential side effects of these medications and instructed to seek medical attention if they experience any adverse effects.

Analgesics are commonly used in dentistry to manage pain and discomfort associated with various dental procedures, such as tooth extractions, root canal treatments, and dental fillings.⁹ However, there are certain contraindications or situations in which the use of analgesics may not be appropriate. Some of the common contraindications for use of analgesics in dentistry include:

Allergy: Patients with a known allergy or hypersensitivity to a specific analgesic or its components should avoid its use. Renal impairment: Patients with impaired kidney function may not be able to eliminate certain analgesics from their body, leading to an increased risk of toxicity. Hepatic impairment: Patients with impaired liver function may have difficulty metabolizing certain analgesics, leading to an increased risk of toxicity.¹⁰ Respiratory depression: Analgesics such as opioids can cause respiratory depression, which can be life-threatening in some cases, especially in patients with pre-existing respiratory conditions. Pregnancy and lactation: Some analgesics may not be safe for use during pregnancy or breastfeeding, as they may affect fetal development or be passed on to the infant through breast milk. Drug interactions: Patients taking certain medications, such as antidepressants or anticoagulants, may experience adverse effects or interactions when taking analgesics. It is important for dental professionals to carefully review a patient's medical history and current medications before prescribing or administering analgesics to ensure their safe and appropriate use.¹¹

3. Role of Antibiotics in Dentistry

Antibiotics are medications that are used to treat bacterial infections. In dentistry, antibiotics are commonly used to treat dental infections, such as periodontitis, dental abscesses, and necrotizing ulcerative gingivitis. Periodontitis is a bacterial infection that affects the gums and the bone that supports the teeth. It can cause gum inflammation, bleeding, and tooth loss. In addition to scaling and root planing, antibiotics may be prescribed to help control the bacterial infection. Antibiotics used to treat periodontitis include tetracyclines, metronidazole, and amoxicillin. Dental abscesses are localized infections that occur around the root of a tooth or in the gum tissue. They can cause severe pain, swelling, and fever. Treatment of dental abscesses typically involves drainage of the pus and the use of antibiotics to control the infection. Antibiotics commonly used to treat dental abscesses include penicillin, amoxicillin, and clindamycin.¹²

Necrotizing ulcerative gingivitis (NUG) is a bacterial infection that affects the gums and the tissues surrounding the teeth. It can cause severe pain, bleeding, and bad breath. Treatment of NUG includes deep cleaning of the affected area and the use of antibiotics to control the bacterial infection. Antibiotics commonly used to treat NUG include metronidazole and amoxicillin. It's important to note that antibiotics should only be prescribed when there is a confirmed bacterial infection that cannot be controlled by other means, such as deep cleaning or root canal therapy. Overuse or misuse of antibiotics can lead to antibiotic resistance, which can make it difficult to treat bacterial infections in the future. Additionally, antibiotics may cause adverse effects, such as allergic reactions, upset

stomach, and diarrhea. Patients should be instructed to take antibiotics as prescribed and to seek medical attention if they experience any adverse effects. Antibiotics should always be prescribed by a healthcare provider after a thorough evaluation of the patient's medical history and symptoms.

3.1. Use and misuse of antibiotics in dentistry

Antibiotics are commonly used in dentistry to treat bacterial infections. However, their use should be judicious to avoid contributing to antibiotic resistance, which is a growing concern worldwide. Antibiotic misuse in dentistry can occur when antibiotics are prescribed unnecessarily, the wrong antibiotic is prescribed, or the patient is not instructed on how to use the medication correctly. One common example of antibiotic misuse in dentistry is the prescription of antibiotics for dental pain or inflammation without a clear indication of bacterial infection. For instance, prescribing antibiotics for routine dental extractions or implant placements is usually not necessary unless there is a specific risk of infection or the patient is at high risk of developing a systemic infection. Another issue is the inappropriate selection of antibiotics. Dentists should consider the patient's medical history, allergies, and potential drug interactions before prescribing an antibiotic. Additionally, dentists should avoid prescribing broad-spectrum antibiotics unnecessarily and should use narrow-spectrum antibiotics when possible. Finally, it is essential to educate patients on the proper use of antibiotics, including the importance of taking the full course of medication as prescribed and not sharing antibiotics with others. Patients should also be advised on potential side effects and what to do if they experience an allergic reaction.

In conclusion, while antibiotics are an essential tool in dentistry, their misuse can have serious consequences, such as promoting antibiotic resistance. Dentists should be cautious when prescribing antibiotics and should follow evidence-based guidelines to ensure optimal patient care.¹³

3.2. Role of local anesthetics in dentistry

Local anesthetics are medications that are used to numb a specific area of the body to provide pain relief during a dental procedure. In dentistry, local anesthetics are commonly used to numb the mouth, gums, and teeth during procedures such as tooth extraction, root canal therapy, and filling cavities. Local anesthetics work by blocking the transmission of pain signals from the nerves in the affected area to the brain. They do not affect consciousness or other areas of the body, which makes them a safe and effective way to manage pain during dental procedures.¹⁴

There are different types of local anesthetics used in dentistry, including lidocaine, articaine, bupivacaine, and mepivacaine. Lidocaine is the most commonly used local

anesthetic in dentistry due to its rapid onset and long duration of action. Articaine is another commonly used local anesthetic that has a faster onset of action and can provide longer-lasting pain relief. Local anesthetics are administered using a syringe, which delivers the medication into the affected area. The medication can take a few minutes to take effect, and patients may feel a tingling or numb sensation in the area. The duration of action of local anesthetics depends on the type and dosage of the medication, as well as the individual patient's metabolism.¹⁵

Local anesthesia is commonly used in dentistry to numb the area where a dental procedure is to be performed. However, there are certain contraindications or situations in which local anesthesia should not be used or used with caution. Some of these contraindications include: Allergy or hypersensitivity to local anesthetics: Patients who have experienced an allergic reaction or have a known hypersensitivity to a specific local anesthetic should avoid using that anesthetic or any related ones. Medical conditions: Patients with certain medical conditions, such as uncontrolled high blood pressure, uncontrolled diabetes, liver disease, or kidney disease, may be at a higher risk of complications from local anesthesia. Drug interactions: Certain medications, such as those used to treat heart conditions, can interact with local anesthetics and cause adverse effects. Infection or inflammation at the site of injection: Local anesthesia should not be used in areas where there is an active infection or inflammation as this may cause further complications. Neurological disorders: Patients with certain neurological disorders, such as multiple sclerosis or Parkinson's disease, may be at a higher risk of complications from local anesthesia. It is important for dentists to take a thorough medical history of their patients and assess any potential risks before administering local anesthesia. In cases where local anesthesia is contraindicated, alternative pain management strategies can be used.¹⁶

3.3. Role of fluorides in dentistry

Fluorides play an important role in maintaining good oral health and preventing tooth decay in dentistry. Fluoride is a mineral that occurs naturally in water, soil, and certain foods. It has been proven to strengthen tooth enamel and make it more resistant to decay. Fluorides are commonly used in dentistry in the following ways:¹⁷ Topical fluoride: Topical fluoride is applied directly to the teeth in the form of gels, foams, or varnishes. It is commonly used in dental offices during routine cleanings and checkups to help prevent tooth decay. Topical fluoride can also be applied at home using fluoride toothpaste or mouthwash. Systemic fluoride: Systemic fluoride is ingested in the form of fluoridated water, dietary supplements, or prescription fluoride tablets. It helps strengthen tooth enamel from within and is especially important for children whose teeth are still

developing. Professional fluoride treatments: Professional fluoride treatments are administered by dental professionals and are more concentrated than over-the-counter fluoride products. These treatments are typically recommended for patients who are at a higher risk for tooth decay, such as those with a history of cavities or those with braces.¹⁸ Fluorides work by strengthening the tooth enamel and making it more resistant to acid attacks from plaque bacteria. They also help reverse early stages of tooth decay by remineralizing weakened areas of the enamel. However, it's important to note that excessive fluoride intake can lead to dental fluorosis, a condition that affects the appearance of the teeth. This condition is characterized by white or brown spots on the teeth and can occur when children consume too much fluoride during the time when their teeth are developing. Dental healthcare providers should always evaluate the patient's fluoride intake and adjust the fluoride treatment accordingly to avoid dental fluorosis.¹⁹

4. Conclusion

It's important for dental healthcare providers to assess each patient's individual needs and medical history to determine the most appropriate pharmacotherapy. They should also educate their patients on the proper use of medications and any potential side effects. By incorporating pharmacotherapy as part of a comprehensive dental care plan, patients can experience a more comfortable and successful dental experience.

5. Source of Funding

None.

6. Conflict of Interest

None.

References

- Jacobs MH. Chemotherapeutics and antibiotics and dentistry. . *Oral Surg Oral Med Oral Pathol*. 1950;3(10):1247–56.
- Hargreaves K, Abbott PV. Drugs for pain management in dentistry. *Aust Dent J*. 2005;50(4):14–22.
- Ahmadi H, Ebrahimi A, Ahmadi F. Antibiotic Therapy in Dentistry. *Int J Dent*. 2021;p. 6667624. doi:10.1155/2021/6667624.
- Lombardi A, Ouanounou A. Fungal infections in dentistry: Clinical presentations, diagnosis, and treatment alternatives. *Oral Surg Oral Med Oral Pathol Oral Radiol*. 2020;130(5):533–46.
- Vallejo AP, García-Pola-Vallejo MJ. Local anesthetics in dentistry. *Med Oral Patol Oral Cir Bucal*. 2004;9:440–3.
- Corcuera-Flores JR, Rangil JS, Soriano AC, Jiménez JL. Current methods of sedation in dental patients - a systematic review of the

- literature. *Med Oral Patol Oral Cir Bucal*. 2016;21(5):579–86.
- Seymour RA. Drug interactions in dentistry. *Dent Update*. 2009;36(8):469–70.
- Roda RP, Bagán JV, Soriano J, Romero G. Use of nonsteroidal antiinflammatory drugs in dental practice. A review. . *Med Oral Patol Oral Cir Bucal*. 2007;12(1):10–8.
- Nagi R, Devi Y, Rakesh BK, Reddy N, Patil SS. Clinical implications of prescribing nonsteroidal anti-inflammatory drugs in oral health care—a review. *Oral Surg Oral Med Oral Pathol Oral Radiol*. 2015;119(3):264–71.
- Hersh EV, Moore PA, Grosser T, Polomano RC, Farrar JT, Saraghi M. Nonsteroidal Anti-Inflammatory Drugs and Opioids in Postsurgical Dental Pain. *J Dent Res*. 2020;99(7):777–86.
- Ahmadi H, Ebrahimi A, Ahmadi F. Antibiotic Therapy in Dentistry. *Int J Dent*. 2021;p. 6667624.
- Oberoi SS, Dhingra C, Sharma G, Sardana D. Antibiotics in dental practice: how justified are we. *Int Dent J*. 2015;65(1):4–10.
- Cope AL, Chestnutt IG. Inappropriate prescribing of antibiotics in primary dental care: reasons and resolutions. *Prim Dent J*. 2014;3(4):33–40.
- Ogle OE, Mahjoubi G. Advances in local anesthesia in dentistry. *Dent Clin North Am*. 2011;55(3):21726685.
- Mathison M, Pepper T. Local Anesthesia Techniques In Dentistry and Oral Surgery. *StatPearls*. 2023;p. 35593805.
- Whelton HP, Spencer AJ, Do LG, Rugg-Gunn AJ. Fluoride Revolution and Dental Caries: Evolution of Policies for Global Use. *J Dent Res*. 2019;98(8):837–46.
- Buzalaf M, Pessan JP, Honório HM, Ten Cate JM. Mechanisms of action of fluoride for caries control. *Monogr Oral Sci*. 2011;22:97–114.
- Pollick H. The Role of Fluoride in the Prevention of Tooth Decay. *Pediatr Clin North Am*. 2018;65(5):923–40.
- Kanduti D, Sterbenk P, Artnik B. Fluoride: a review of use and effects on health. *Mater Sociomed*. 2016;28(2):133–40.

Author biography

Shailesh Shenoy, Associate Professor

Zameer Pasha, Consultant in Oral and Maxillofacial Surgery

Suresh Rajendiran, Associate Professor

Syed Owais Ahmed, Assistant Professor

Krishan Kant, Resident

Akshaya Narayan Shetti, 0000-0002-4688-8071
 <https://orcid.org/0000-0002-4688-8071>

Cite this article: Shenoy S, Pasha Z, Rajendiran S, Ahmed SO, Kant K, Shetti AN. The intersection of pharmacology and dentistry: A comprehensive overview of pharmacotherapeutics. *Int J Pharm Chem Anal* 2023;10(2):66-69.